


We claim:

A carboxylic acid derivative of the formula I



where R is formyl, tetrazole ~~[sic]~~, nitrile ~~[sic]~~, a COOH group or a radical which can be hydrolyzed to COOH, and the other substituents have the following meanings:

R² hydrogen, hydroxyl, NH₂, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio;

X nitrogen or CR¹⁴ where R¹⁴ is hydrogen or C₁₋₅-alkyl, or CR¹⁴ forms together with CR³ a 5- or 6-membered alkylene or alkenylene ring which can be substituted by one or two C₁₋₄-alkyl groups and in which in each case a methylene group can be replaced by oxygen, sulfur, -NH or -NC₁₋₄-alkyl;

R³ hydrogen, hydroxyl, NH₂, NH(C₁-C₄-Alkyl), N(C₁-C₄-alkyl)₂, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, -NH-O-C₁₋₄-alkyl, C₁-C₄-alkylthio or CR³ is linked to CR¹⁴ as indicated above to give a 5- or 6-membered ring;

R⁴ and R⁵ (which can be identical or different):

phenyl or naphthyl, which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, amino, C₁-C₄-alkylamino or C₁-C₄-di-alkylamino; or

phenyl or naphthyl, which are connected together in the ortho positions via a direct linkage, a methylene, ethylene or ethenylene group, an oxygen or sulfur atom or an SO₂, NH or N-alkyl group

44

202506960329

or C₃-C₇-cycloalkyl;

B1
cont

5 R⁶ hydrogen, C₁-C₈-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or C₃-C₈-cycloalkyl, where each of these radicals can be substituted one or more times by: halogen, nitro, cyano, C₁-C₄-alkoxy, C₃-C₆-alkenyloxy, C₃-C₆-alkynyloxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxy-carbonyl, C₃-8-alkylcarbonylalkyl, C₁-C₄-alkylamino, di-C₁-C₄-alkylamino, phenyl or phenyl or phenoxy which is substituted one or more times, eg. one to three times, by

10 halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio;

phenyl or naphthyl, each of which can be substituted by one

15 or more of the following radicals: halogen, nitro, cyano, hydroxyl, amino, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, C₁-C₄-dialkylamino or dioxomethylene ~~[sic]~~ or dioxoethylene ~~[sic]~~;

20 a five- or six-membered heteroaromatic moiety containing one to three nitrogen atoms and/or one sulfur or oxygen atom, which can carry one to four halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl,

25 C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, phenyl, phenoxy or phenylcarbonyl, it being possible for the phenyl radicals in turn to carry one to five halogen atoms and/or one to three of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or

30 C₁-C₄-alkylthio;

with the proviso that R⁶ can be hydrogen only when Z is not a single bond;

35 Y sulfur or oxygen or a single bond;

2 Z sulfur, oxygen, -SO-, -SO₂- ~~or a single bond.~~

40

add

45